

Lamination of the outer plexiform layer in optic atrophy caused by dominant *WFS1* mutations

Majander A¹, Bitner-Glindzicz M², Chan CM¹, Duncan HJ³, Chinnery PF^{4,5,6}, Subash M¹, Keane PA¹, Webster AR¹, Moore AT^{1,7}, Michaelides M¹, Yu-Wai-Man P^{1,3,4}

¹Moorfields Eye Hospital and UCL Institute of Ophthalmology, London, UK.

²Great Ormond Street Hospital, Great Ormond Street, London, UK.

³Newcastle Eye Centre, Royal Victoria Infirmary, Newcastle upon Tyne, UK.

⁴Wellcome Trust Centre for Mitochondrial Research, Institute of Genetic Medicine, Newcastle University, Newcastle upon Tyne, UK.

⁵Medical Research Council Mitochondrial Biology Unit, Cambridge, UK

⁶Department of Clinical Neurosciences, School of Clinical Medicine, University of Cambridge, UK

⁷ Ophthalmology Department, UCSF School of Medicine, San Francisco, CA

Corresponding Authors

Dr Anna Majander

UCL Institute of Ophthalmology and Moorfields Eye Hospital

162 City Road, London, EC1V 2PD, United Kingdom

Tel: +44-0207 566 2265

E-mail: a.majander@ucl.ac.uk

Dr Patrick Yu-Wai-Man

Wellcome Trust Centre for Mitochondrial Research

Institute of Genetic Medicine, Newcastle University

Newcastle upon Tyne, NE1 3BZ, United Kingdom

Tel: +44-191-241-8854, Fax: +44-191-241-8666

E-mail: Patrick.Yu-Wai-Man@ncl.ac.uk

Financial Support:

This research has been supported by the National Institute for Health Research Rare Diseases Translational Research Collaboration (NIHR RD-TRC) and the National Institute for Health Research Biomedical Research Centre at Moorfields Eye Hospital National Health Service Foundation Trust and UCL Institute of Ophthalmology. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. AM also receives funding from Suomen Silmätutkimusseura ry:n Apurahasäätiö (Finland). MM also receives funding from Fight For Sight (UK), Moorfields Eye Hospital Special Trustees, Moorfields Eye Charity, the Foundation Fighting Blindness (USA), Retinitis Pigmentosa Fighting Blindness, and The Wellcome Trust [099173/Z/12/Z]. MM is supported by an FFB Career Development Award. PAK is a NIHR Clinician Scientist. PFC is a Wellcome Trust Senior Fellow in Clinical Science and a UK NIHR Senior Investigator who also receives funding from the MRC (UK), Parkinson's UK, and the UK NIHR Biomedical Research Centre for Ageing and Age-related disease award to the Newcastle upon Tyne Hospitals NHS Foundation Trust. PYWM is a Medical Research Council (MRC, UK) Clinician Scientist. PYWM also receives funding from Fight for Sight (UK) and the UK National Institute of Health Research (NIHR) as part of the Rare Diseases Translational Research Collaboration. The funding organizations had no role in the design or conduct of this research.

Disclosures

All the listed authors in this manuscript report no relevant financial disclosures or conflicts of interest.

Running title

OPL lamination in optic atrophy secondary to dominant *WFS1* mutations

Keywords

Autosomal dominant optic atrophy; DIDMOAD; optical coherence tomography; WFS1; Wolfram syndrome

Address for reprints

Dr Patrick Yu-Wai-Man

Wellcome Trust Centre for Mitochondrial Research

Institute of Genetic Medicine, Newcastle University

Newcastle upon Tyne, NE1 3BZ, United Kingdom

Tel: +44-191-241-8854, Fax: +44-191-241-8666

E-mail:Patrick.Yu-Wai-Man@ncl.ac.uk

Word count: 1,012

Title: 77 characters (no spaces)

One figure: Figure 1

2 online supplement tables: Table 1 and Table 2

1 Wolfram syndrome, or DIDMOAD (diabetes insipidus, diabetes mellitus, optic atrophy (OA), and
2 deafness), is a neurodegenerative disorder with heterogeneous clinical manifestations caused by
3 homozygous or compound heterozygous recessive mutations in the *WFS1* gene (OMIM 606201).¹
4 More recently, the phenotypic spectrum has expanded to include patients with dominant
5 inheritance and limited clinical features, in particular OA in association with diabetes mellitus and/or
6 sensorineural deafness.¹ *WFS1* encodes for an endoplasmic reticulum transmembrane protein,
7 Wolframin, which is highly expressed in retinal tissues, including retinal ganglion cells, the
8 photoreceptor inner segments, and the inner nuclear layer (INL) of the human eye, and mouse
9 Müller cells.^{2,3} Thinning of the peripapillary nerve fibre and macular retinal ganglion cell layers are
10 typical features of *WFS1*-related optic atrophy, but careful genotype-phenotype correlations have
11 not yet been established. Here, we report on a comprehensive macular OCT imaging study of 14
12 patients with OA secondary to mutations in *WFS1* and the identification of a distinct outer plexiform
13 layer (OPL) abnormality, which associates exclusively with dominant missense *WFS1* mutations
14 (**Table 1**, available at www.aaojournal.org).

15 Averaged B-scans of 26 eyes from 14 patients with confirmed pathogenic *WFS1* mutations
16 were retrospectively reviewed from our Spectralis™ (Heidelberg Engineering Ltd., Heidelberg,
17 Germany) databases. A prominent feature in the macular SD-OCT images of 15 eyes from 8 patients
18 harbouring dominant *WFS1* mutations (Patients 1-8) was an abnormal reflectivity of the OPL, which
19 was not observed in 11 eyes from 6 patients with recessive *WFS1* mutations (**Figure 1** and **Table 1**,
20 available at www.aaojournal.org). The OPL was comprised of three distinct laminae: an innermost
21 highly reflective lamina; a middle non-reflective lamina; and an outermost highly reflective lamina.
22 To a certain extent, OPL thickness varied depending on the angle between the OCT laser beam and
23 the retinal layers. Beam positions that were more perpendicular to the Henle's fibre tracks resulted
24 in a thicker OPL compared with beam positions that were parallel to these tracks, as illustrated by

the left eye of Patient 5 (**Figure 1**). However, in several eyes, an abnormally thick OPL was visualized even with straight OCT beam positions, as for Patient 6 (**Figure 1**). In the majority of patients, the middle OPL lamina formed a nearly confluent ring centred at the fovea and extending 1-1.5 mm from the foveolar centre up to the optic disc. Perifoveal volumetric retinal scans were obtained for 22 eyes from 12 patients in our *WFS1* cohort. Automated segmentation and retinal layer thickness analysis were carried out using the automated retinal Heidelberg Engineering segmentation tool included in the Spectralis Glaucoma Module software (version 6.0) after manual confirmation of each layer (**Table 2**, available at www.aaojournal.org). Marked thinning of the GCL-IPL complex was observed consistent with OA. All 11 eyes with abnormal OPL lamination had significantly thicker OPL and, more importantly, combined OPL and ONL complex compared with control eyes, or the eyes of patients without OPL lamination. Compared with control eyes, INL thickness was increased in 4 eyes (Patients 2, 5, 8) with INL cystoid changes (**Figure 1**), but also in 5 other eyes with OPL lamination and in 3 eyes without OPL abnormality (**Table 2**, available at www.aaojournal.org). Our study had ethical and institutional approval and its design complied with the Declaration of Helsinki.

We have identified a previously unreported macular OCT feature in patients with Wolfram-related optic atrophy that is associated with dominant, but not recessive, *WFS1* mutations. The lamination of the OPL was characterised by three distinct sublayers: (1) an innermost highly-reflective lamina, similar to what is frequently identified as the OPL in normal eyes; (2) a non-reflective cleft-like middle lamina; and (3) an outermost highly-reflective lamina with characteristics of Henle's fibre layer. Our retrospective study did not enable a systematic evaluation of the effect of angle changes on the OPL reflectivity. Further work is needed to clarify whether the outermost lamina has truly abnormal reflectivity and whether the middle lamina represents an additional spatial structure or merely abnormal OCT reflectivity. However, as the thickness of combined OPL and ONL complex should not be sensitive to OCT angle changes, its significantly increased thickness

in eyes with abnormal OPL lamination suggests that an additional space could indeed account for the laminated OPL structure observed in this patient subgroup with dominant *WFS1* mutations.

Taking into consideration the spatial relationships between the three OPL laminae and previously published histological studies,⁴ we propose that the middle cleft-like lamina is located between the synaptic and pedicles sublayers of the OPL. This region has also been shown to possess a particular retinal architecture with bending Müller cells.⁵ Wolframin is highly expressed in Müller cells and this cell type has been implicated in the development of OPL and INL edema, and changes within Henle's fibre layer.^{4,5} The Müller cell is, therefore, an attractive candidate for the development of the observed OPL lamination.

Interestingly, OPL lamination was only associated with dominant missense *WFS1* mutations. It was not identified in any of the 6 patients harbouring recessive homozygous or compound heterozygous *WFS1* mutations. This distinct OPL abnormality has not previously been reported in the context of optic atrophy and it could represent a specific deleterious effect of dominant *WFS1* mutations. The missense variants identified in our patients are predicted to result in only a minor reduction in wolframin protein level.¹ Depletion of the native wild-type wolframin protein, which is thought to be the major pathophysiological mechanism in recessive Wolfram syndrome, is therefore unlikely to be implicated in the aetiology of the OPL lamination. Instead, dominant missense *WFS1* mutations could result in a dysfunctional aberrant wolframin protein.

WFS1 is a highly polymorphic gene and determining the mode of inheritance or the pathological significance of a specific *WFS1* variant is not always straightforward, especially in singleton cases with no access to other family members. Our study has revealed an interesting association between dominant missense *WFS1* mutations and distinct OPL lamination on SD-OCT, which was not observed in patients with recessive *WFS1* mutations. Future work in other independent patient cohorts with Wolfram syndrome will confirm whether this OCT feature could be used to distinguish between dominant and recessive forms of the disease.

References

- 1 de Heredia ML, Clèries R, Nunes V. Genotypic classification of patients with Wolfram syndrome: insights into the natural history of the disease and correlation with phenotype. *Genet Med* 2013;15:497-506.
- 2 Schmidt-Kastner R, Kreczmanski P, Preising M, et al. Expression of the diabetes risk gene wolframin (WFS1) in the human retina. *Exp Eye Res* 2009;89:568-74.
- 3 Kawano J, Tanizawa Y, Shinoda K. Wolfram syndrome 1 (Wfs1) gene expression in the normal mouse visual system. *J Comp Neurol* 2008;510:1-23
- 4 Curcio CA, Messinger JD, Sloan KR, et al. Human chorioretinal layer thicknesses measured in macula-wide, high-resolution histologic sections. *Invest Ophthalmol Vis Sci* 2011;52:3943-54.
- 5 Matet A, Savastano MC, Rispoli M, et al. En face optical coherence tomography of foveal microstructure in full-thickness macular hole: A model to study perifoveal Müller cells. *Am J Ophthalmol* 2015;159:1142–51.

90 **A figure legend:**

91 **Figure 1.** Macular SD-OCT images of Patient 1 showing outer plexiform layer (OPL) lamination and of
92 Patient 13 without this defect (upper panel). The lower panels show OCT images of Patients 2-8 with
93 OPL lamination, with the right eye on the left and the left eye on the right. The localization of the
94 layers that were not imaged through the optical axis are indicated within the inserts. The yellow
95 arrows indicate the cleft-like middle structure of the OPL. The red arrows highlight areas of cystoid
96 inner nuclear layer (INL) edema.




Table 1. *WFS1* genotype and clinical characteristics of the patients in our study cohort.

Patient	Sex	Inheritance (Family)	<i>WFS1</i> mutation(s)*	Extraocular findings	OA age (y)**	BCVA (logMAR)		Macular SD-OCT	
						RE	LE	Age (y)**	OPL cleft***
1	F	AD (1)	c.2390A>T: p.Asp797Val	SNHL, DM2	45	1.10	NI	69	+
2	F	AD (2)	c.2051C>T: p.Ala684Val	SNHL, epilepsy, hypertension	5	0.60	0.60	46	+
3	M	AD (2)	c.2051C>T: p.Ala684Val	SNHL	23	0.48	0.60	23	+
4	F	AD (3)	c.968A>G: p.His323Arg	SNHL	U	0.30	0.43	44	+
5	F	AD (3)	c.968A>G: p.His323Arg	SNHL	4	0.00	0.00	8	+
6	M	<i>de novo</i> (4)	c.937C>T: p.His313Tyr c.-170-9_14dupTGCCCC	SNHL, DM1, short stature	7	0.10	0.20	10	+
7	F	AD (5)	c.2051C>T: p.Ala684Val	SNHL	78	1.98	1.98	78	+
8	F	AD (6)	c.2161A>T: p.Asn721Tyr	SNHL	50	1.98	1.98	51	+
9	M	AR (7)	c.505G>A: p.Glu169Lys c.874C>A: p.Pro292Thr	DM1, DI, NB	9	2.28	2.28	45	-
10	M	AR (7)	c.505G>A: p.Glu169Lys c.874C>A: p.Pro292Thr	DM1, DI, NB	U	1.30	NI	46	-
11	F	AR (8)	c.2648_2651delTCTT: p.Phe883SerfsX68 c.1597C>T: p.Pro533Ser	DI UC	23	0.60	0.60	44	-
12	F	AR (8)	c.2648_2651delTCTT: p.Phe883SerfsX68 c.1597C>T: p.Pro533Ser	DI UC	37	0.78	0.78	52	-
13	F	AR (9)	c.2643_2644delC: p.Phe883LeufsX56 (homozygous)	SNHL, DM1, DI PS, NEU	15	1.30	1.30	25	-
14	F	AR (10)	c.409_424dup1: p.Val142GlyfsX10 c.2262_2263delCT: p.Cys755SerfsX3	SNHL, DM1	31	1.00	1.00	35	-

* *WFS1* genetic testing was conducted by the West Midlands Regional Genetic Laboratory, Birmingham, UK.

** Age at the time of diagnosis of optic atrophy (OA) and when macula SD-OCT imaging was performed. Images were retrieved from the Spectralis™ (Heidelberg Engineering Ltd., Heidelberg, Germany) databases of Moorfields Eye Hospital, London, UK and the Newcastle Eye Centre, Royal Victoria Infirmary, Newcastle upon Tyne, UK. Our study had local ethical and institutional approval, and its design complied with the Declaration of Helsinki.

***The presence (+) or absence (-) of an outer plexiform layer (OPL) lamination is indicated.

Abbreviations: DI, Diabetes insipidus; DM1, Diabetes mellitus type 1; DM2, Diabetes mellitus type 2; LE, left eye; NB, Neurogenic bladder; NEU, intermittent myoclonus, occasional balance problem, postural hypotension; NI, not included into the study; NP, not performed; OA; optic atrophy; OPL; outer plexiform layer; PS, bipolar affective disorder RE, right eye; RNFL, peripapillary retinal nerve fiber layer thickness; SNHL, sensorineural hearing loss; U, Unknown; UC, urinary and faecal incontinence.